

MEDICATION GUIDE
Eszopiclone (ES-zoe-PIK-lone)
Tablets USP ©
Rx only

Read the Medication Guide that comes with eszopiclone tablets before you start taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your doctor about your medical condition or treatment.

What is the most important information I should know about eszopiclone tablets?

- Do not take more eszopiclone tablets than prescribed.
- Do not take eszopiclone tablets unless you are able to stay in bed a full night (7 to 8 hours) before you must be active again.
- Take eszopiclone tablets right before you get in bed, not sooner.

Eszopiclone tablets may cause serious side effects that you may not know are happening to you. These side effects include:

- sleepiness during the day
- not thinking clearly
- act strangely, confused, or upset
- “sleep-walking” or doing other activities when you are asleep like:
 - eating
 - talking
 - having sex
 - driving a car

Call your healthcare provider right away if you find out that you have done any of the above activities after taking eszopiclone tablets.

The morning after you take eszopiclone tablets your ability to drive safely and think clearly may be decreased.

Do not take eszopiclone tablets if you:

- drank alcohol that evening or before bed
- take other medicines that can make you sleepy. Talk to your doctor about all of your medicines. Your doctor will tell you if you can take eszopiclone tablets with your other medicines.
- cannot get a full night’s sleep

WHAT IS ESZOPICLONE TABLET?

Eszopiclone tablet is a sedative-hypnotic (sleep) medicine. Eszopiclone tablet is used in adults for the treatment of a sleep problem called insomnia. Symptoms of insomnia include:

- trouble falling asleep
- waking up often during the night

Eszopiclone tablets are not for children.

Eszopiclone tablet is a federally controlled substance (C-IV) because it can be abused or lead to dependence. Keep eszopiclone tablets in a safe place to prevent misuse and abuse. Selling or giving away eszopiclone tablets may harm others, and is against the law. Tell your doctor if you have ever abused or been dependent on alcohol, prescription medicines or street drugs.

Who should not take eszopiclone tablets?

Do not take eszopiclone tablets if you are allergic to anything in it.

See the end of this Medication Guide for a complete list of ingredients in eszopiclone tablets.

Eszopiclone tablets may not be right for you. Before starting eszopiclone tablets, tell your doctor about all of your health conditions, including if you:

- have a history of depression, mental illness, or suicidal thoughts
- have a history of drug or alcohol abuse or addiction
- have liver disease
- are pregnant, planning to become pregnant, or breastfeeding

Tell your doctor about all of the medicines you take including prescription and nonprescription medicines, vitamins and herbal supplements. Medicines can interact with each other, sometimes causing serious side effects. **Do not take eszopiclone tablets with other medicines that can make you sleepy.**

Know the medicines you take. Keep a list of your medicines with you to show your doctor and pharmacist each time you get a new medicine.

How should I take eszopiclone tablets?

- Take eszopiclone tablets exactly as prescribed. Do not take more eszopiclone tablets than prescribed for you.
- **Take eszopiclone tablets right before you get into bed.**
- Do not take eszopiclone tablets with or right after a meal.
- **Do not take eszopiclone tablets unless you are able to get a full night’s sleep before you must be active again.**
- **Call your doctor if your insomnia worsens or is not better within 7 to 10 days.** This may mean that there is another condition causing your sleep problems.
- If you take too much eszopiclone tablets or overdose, call your doctor or poison control center right away, or get emergency treatment.

SAP Code: 60100001945
259644 Eszopiclone Tabs_M.G v_2.pdf

Issued For Final Production :-	
Issued For Re-Approval :-	
Issued For Approval :-	22-Feb-19 10:11:46
TEMPLE Packaging Pvt. Ltd. Tel:- 28476601, Email:- info@templepackaging.co.in	
Customer :- LUPIN LTD.	SAP Code :- 60100001945
Location :- Goa	File No. :- 56181
Product Name :- MED GUIDE ESZOPICLONE TABS USP US R4	Artwork Sr. No. :- 169/02/19
Product Code :- 259644	Customer A/W No. :- NA
Version No. :- 02	Old File No. :- 46240
Date :- 22-Feb-19	Old Product Code :- 250806
Artwork Status :- FINAL	Barcode :- NA
Open Size :- 250X140 MM	Pharmacode :- NA
Folding Size :- 24Pgs/ booklet	Perforation :- NA
Substrate :- 60 GSM MAPLITHO PAPER	Gluing :- NA
Cover Page :- NA	Font Name :- Helvetica Condensed
	Font Size :- 10pt.
Remark :- NA	
Printing Colours 1	

K

Note - This approval will be considered for final printing. Please recheck for corrections indicated earlier, in this proof also.			
Prepared by : 22-Feb-19 sanjay	Checked by :	Q.A. Checked :	Approved by :
CUSTOMER APPROVAL			
Checked By		Approved By	
Packaging Development	Production	Regulatory Affairs	Quality Assurance

What are the possible side effects of eszopiclone tablets?

Possible serious side effects of eszopiclone tablets include:

- **getting out of bed while not being fully awake and do an activity that you do not know you are doing.** (See “What is the most important information I should know about eszopiclone tablets?”)
- **abnormal thoughts and behavior.** Symptoms include more outgoing or aggressive behavior than normal, confusion, agitation, hallucinations, worsening of depression, and suicidal thoughts or actions.
- **memory loss**
- **anxiety**
- **severe allergic reactions.** Symptoms include swelling of the tongue or throat, trouble breathing, and nausea and vomiting. Get emergency medical help if you get these symptoms after taking eszopiclone tablets.

Call your doctor right away if you have any of the above side effects or any other side effects that worry you while using eszopiclone tablets.

The most common side effects of eszopiclone tablets are:

- unpleasant taste in mouth, dry mouth
- drowsiness
- dizziness
- headache
- symptoms of the common cold
- You may still feel drowsy the next day after taking eszopiclone tablets. **Do not drive or do other dangerous activities after taking eszopiclone tablets until you feel fully awake.**

These are not all the side effects of eszopiclone tablets. Ask your doctor or pharmacist for more information. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Lupin Pharmaceuticals, Inc. at 1-800-399-2561.

How should I Store eszopiclone tablets?

- Store eszopiclone tablets at room temperature, between 59° F to 86° F (15° C to 30° C).
- Do not use eszopiclone tablets after the expiration date.
- **Keep eszopiclone tablets and all medicines out of reach of children.**

General Information about eszopiclone tablets

- Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide.

- Do not use eszopiclone tablets for a condition for which it was not prescribed.
- Do not share eszopiclone tablets with other people, even if you think they have the same symptoms that you have. It may harm them and is against the law.

This Medication Guide summarizes the most important information about eszopiclone tablets. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about eszopiclone tablets that is written for healthcare professionals.

- To report side effects, call 1-800-399-2561.

What are the ingredients in eszopiclone tablets?

Active Ingredient: eszopiclone

Inactive Ingredients: colloidal silicon dioxide, croscarmellose sodium, dibasic calcium phosphate dihydrate, hypromellose, lactose monohydrate, polyethylene glycol, povidone, pregelatinized starch, magnesium stearate and titanium dioxide. In addition, both the 1 mg and 3 mg tablets contain FD&C Blue #2.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Manufactured for:

Lupin Pharmaceuticals, Inc.
Baltimore, Maryland 21202
United States

Manufactured by:

Lupin Limited
Goa 403 722
INDIA

Revised: March 6, 2019

ID#: 259644

**SAP Code: 60100001945
259644 Eszopiclone Tabs_M.G v_2.pdf**

Issued For Final Production :-	
Issued For Re-Approval :-	
Issued For Approval :-	22-Feb-19 10:11:46
TEMPLE Packaging Pvt. Ltd. Tel:- 28476601, Email:- info@templepackaging.co.in	
Customer :- LUPIN LTD.	SAP Code :- 60100001945
Location :- Goa	File No. :- 56181
Product Name :- MED GUIDE ESZOPICLONE TABS USP US R4	Artwork Sr. No. :- 169/02/19
Product Code :- 259644	Customer A/W No. :- NA
Version No. :- 02	Old File No. :- 46240
Date :- 22-Feb-19	Old Product Code :- 250806
Artwork Status :- FINAL	Barcode :- NA
Open Size :- 250X140 MM	Pharmacode :- NA
Folding Size :- 24Pgs/ booklet	Perforation :- NA
Substrate :- 60 GSM MAPLITHO PAPER	Gluing :- NA
Cover Page :- NA	Font Name :- Helvetica Condensed
	Font Size :- 10pt.
Remark :- NA	
Printing Colours	1

K

Note - This approval will be considered for final printing. Please recheck for corrections indicated earlier, in this proof also.			
Prepared by : 22-Feb-19 sanjay	Checked by :	Q.A. Checked :	Approved by :
CUSTOMER APPROVAL			
Checked By		Approved By	
Packaging Development	Production	Regulatory Affairs	Quality Assurance